

SU-4240-635
 hep-ph/9606331
 June, 1996

Pi Pi scattering in an effective chiral Lagrangian*

Masayasu HARADA[†]

*Department of Physics, Syracuse University
 Syracuse, NY 13244-1130, USA*

Abstract

In this talk I summarize a recently proposed mechanism to understand $\pi\pi$ scattering to 1 GeV. The model is motivated by the $1/N_C$ expansion to QCD, and includes a current algebra contact term and resonant pole exchanges. Chiral symmetry plays an important role in restricting the form of the interactions. The existence of a broad low energy scalar (σ) is indicated.

1 Introduction

In this talk I will show the main mechanism studied in Ref. 1, where a simple model of $\pi\pi$ scattering is presented.

The $\pi\pi$ scattering has been studied as an important test of the strong interaction. Now QCD is known to be the fundamental theory of the strong interaction. However, it is very difficult to reproduce the experimental data directly from QCD. One clue is given by the structure of the chiral symmetry, which approximately exists in the QCD Lagrangian and is broken by the strong interaction of QCD. Another clue is given by the $1/N_C$ expansion to QCD. In the large N_C limit, QCD becomes a theory of weakly interacting mesons, and the $\pi\pi$ scattering is expressed as an infinite sum of tree diagrams of mesons.[2]

The experimental data in the low energy region near $\pi\pi$ threshold can be reproduced by using the information from chiral symmetry. This situation is easily understood by using a chiral Lagrangian which includes pions only. In addition, by including the higher derivative terms together with one-loop effects, the applicable energy region is enlarged. This systematic low energy expansion is called the chiral perturbation theory.[3] Using the chiral perturbation theory, we can easily study the result from the chiral symmetry systematically.

* Talk given at Montreal-Rochester-Syracuse-Toronto meeting (May 9–10, 1996).

[†] e-mail: mharada@npac.syr.edu

In the higher energy region, however, the one-loop amplitude of chiral perturbation theory violates the unitarity bound around 400 – 500 MeV in the $I = 0$ S -channel.[4] For the P -wave amplitude, we have the ρ meson, and chiral perturbation theory may break down at the resonance position. The explicit inclusion of resonances in the high energy region easily reproduces the amplitude, of course.

When we apply the large N_C argument to the practical $\pi\pi$ scattering, we cannot actually include an infinite number of resonances. Moreover, the forms of interactions are not fully determined in the large N_C limit. Nevertheless, some encouraging features were previously found in an approach which truncated the particles appearing in the effective Lagrangian to those with masses up to an energy slightly greater than the range of interest.[5] Moreover, the chiral symmetry played an important role to restrict the form of interaction, i.e., the effective Lagrangian was constructed by using the information of chiral symmetry. This seems reasonable phenomenologically and is what one usually does in setting up an effective Lagrangian. In Ref. 5 this Lagrangian provided, as a starting point, a contact term which described the threshold region. However the usual observation was made that the real part of the $I = 0$, $J = 0$ partial wave amplitude quite soon violated the unitarity bound $|R_0^0| \leq 1/2$ rather severely. The inclusion of the contribution coming from the ρ meson exchange was observed to greatly improve, although not completely solve, this problem. These results are shown explicitly in Fig. 1 and provide some encouragement for the possible success of a truncation scheme.

In this talk I concentrate on the energy region below 1 GeV. For the resonances lighter than 1 GeV, ρ and $f_0(980)$ are listed in the particle data group (PDG) list[6] (see Table 1). However, the width of $f_0(980)$ is not well determined. Moreover, the existence of a light scalar σ is suggested by several authors.[7] Here I will determine these resonance parameters by fitting to the $I = 0$ S -wave $\pi\pi$ scattering amplitude.




Figure 1: Predicted curves for R_0^0 . The solid line shows the *current algebra* + ρ result for R_0^0 . The dashed line shows the *current algebra* result alone.

	$I^G(J^{PC})$	$M(\text{MeV})$	$\Gamma(\text{MeV})$
$\sigma(?)$	$0^+(0^{++})$	—	—
$\rho(770)$	$1^+(1^{--})$	769.9	151.2
$f_0(980)$	$0^+(0^{++})$	980	40–400

Table 1: Resonances included in the $\pi\pi \rightarrow \pi\pi$ channel as listed in the PDG. Note that the σ is not present in the PDG and is not being described exactly as a *Breit-Wigner* shape.

This paper is organized as follows. In section 2 I will summarize the resonance model. Section 3 is a main part of this talk, where I will show how to regularize the amplitude, and fit the resonance parameters to the experimental data of the $I = 0$, $J = 0$ partial wave amplitude. Finally, a summary is given in section 4.

2 Resonance Model

In this section I will show how to include higher resonances into the effective chiral Lagrangian.

The most important particle in this energy region is the vector meson. There exist several ways to include the vector meson field into the effective chiral Lagrangian. Here I will include the vector meson as a gauge field of chiral symmetry[8], which is equivalent to the hidden local gauge method (See, for a review, Ref. 9.) at tree level.

Let us start with the non-linear realization of the chiral $U(3)_L \times U(3)_R$ symmetry. The basic quantity is a 3×3 matrix U , which transforms as

$$U \rightarrow U_L U U_R^\dagger, \quad (1)$$

where $U_{L,R} \in U(3)_{L,R}$. This U is parameterized by the pseudoscalar ϕ as

$$U = \xi^2, \quad \xi = e^{2i\phi/F_\pi}, \quad (2)$$

where F_π is a pion decay constant ($F_\pi \simeq 131 \text{ MeV}$). Under the chiral transformation Eq. (1), ξ transforms non-linearly:

$$\xi \rightarrow U_L \xi K^\dagger(\phi, U_L, U_R) = K(\phi, U_L, U_R) \xi U_R^\dagger. \quad (3)$$

The vector meson nonet ρ_μ is introduced as a *gauge field* [8] which transforms as

$$\rho_\mu \rightarrow K \rho_\mu K^\dagger + \frac{i}{\bar{g}} K \partial_\mu K^\dagger, \quad (4)$$

where $\tilde{g}$ is a *gauge coupling constant*. It is convenient to define

$$p_\mu, v_\mu \equiv \frac{i}{2} \left(\xi \partial_\mu \xi^\dagger \mp \xi^\dagger \partial_\mu \xi \right) , \quad (5)$$

which transform as

$$p_\mu \rightarrow K p_\mu K^\dagger , \quad v_\mu \rightarrow K v_\mu K^\dagger + i K \partial_\mu K^\dagger . \quad (6)$$

Using the above quantities we construct the chiral Lagrangian including both pseudoscalar and vector mesons:

$$\mathcal{L} = -\frac{1}{2} m_v^2 \text{Tr} \left[(\rho_\mu - v_\mu / \tilde{g})^2 \right] - \frac{F_\pi^2}{2} \text{Tr} [p_\mu p_\mu] - \frac{1}{4} \text{Tr} [F_{\mu\nu}(\rho) F_{\mu\nu}(\rho)] , \quad (7)$$

where $F_{\mu\nu} = \partial_\mu \rho_\nu - \partial_\nu \rho_\mu - i \tilde{g} [\rho_\mu, \rho_\nu]$ is a *gauge field strength* of vector mesons.[‡]

We next introduce scalar resonances into the Lagrangian. We write the interaction between the scalar nonet field S and pseudoscalar mesons. Under the chiral transformation, this S transforms as $S \rightarrow K S K^\dagger$. A possible form which includes the minimum number of derivatives is proportional to $\text{Tr} [S p_\mu p_\mu]$. The coupling of a physical isosinglet field to two pions is then described by[§]

$$\mathcal{L}_\sigma = -\frac{\gamma_0}{\sqrt{2}} \sigma \partial_\mu \vec{\pi} \cdot \partial_\mu \vec{\pi} . \quad (8)$$

Here we should note that the chiral symmetry requires derivative-type interactions between the scalar field and pseudoscalar mesons.

3 Fit to $\pi\pi$ scattering to 1 GeV

In this section, I will calculate the S -wave $\pi\pi$ scattering amplitude by including resonances as explained in the previous section.

The most problematic feature involved in comparing the leading $1/N_C$ amplitude with experiment is that it does not satisfy unitarity. Since the mesons have no width in the large N_C limit, the amplitude diverges at the resonance position. Thus in order to compare the $1/N_C$ amplitude with experiment we need to regularize the resonance contribution. The

[‡] In actual fit in the next section, we include the pion mass term caused by the explicit chiral symmetry breaking quark mass term. The inclusion of SU(3) symmetry breaking terms in the above Lagrangian is summarized, for example, in Refs. 10 and 11.

[§] The $f_0(980)$ interaction has the same form.

ordinary narrow resonances such as ρ meson are regularized by including the width in the denominator of the propagator:

$$\frac{M\Gamma}{M^2 - s - iM\Gamma} . \quad (9)$$

This is only valid for a narrow resonance in a region where the *background* is negligible. Note that the width in the denominator is related to the coupling constant.

For a very broad resonance there is no guarantee that such a form is correct. Actually, in Ref. 5 it was found necessary to include a rather broad low lying scalar resonance (denoted $\sigma(550)$) to avoid violating the unitarity bound. A suitable form turned out to be of the type

$$\frac{MG}{M^2 - s - iMG'} , \quad (10)$$

where the parameter G' was introduced to regularize the propagator. The important point is that the parameter G' is a free parameter which is not related to the coupling constant.

Even if the resonance is narrow, the effect of the background may be rather important. This seems to be true for the case of the $f_0(980)$. Demanding local unitarity in this case yields a partial wave amplitude of the well known form:[12]

$$\frac{e^{2i\delta} M\Gamma}{M^2 - s - iM\Gamma} + e^{i\delta} \sin \delta , \quad (11)$$

where δ is a background phase (assumed to be slowly varying). We will adopt a point of view in which this form is regarded as a kind of regularization of our model. Of course, non zero δ represents a rescattering effect which is of higher order in $1/N_C$. The quantity $e^{2i\delta}$, taking $\delta = \text{constant}$, can be incorporated into the squared coupling constant connecting the resonance to two pions. In this way, crossing symmetry can be preserved. The non-pole background term in Eq. (11) and hence δ is to be predicted by the other pieces in the effective Lagrangian.

Another point which must be addressed in comparing the leading $1/N_C$ amplitude with experiment is that it is purely real away from the singularities. The regularizations mentioned above do introduce some imaginary pieces but these are clearly more model dependent. Thus it seems reasonable to compare the real part of our predicted amplitude with the real part of the experimental amplitude. Note that the difficulties mentioned above arise only for the direct channel poles; the crossed channel poles and contact terms will give purely real finite contributions.

Let us start from the *current algebra* + ρ contribution. The predicted curve is shown in Fig. 1 in the Introduction. As argued there, although the introduction of ρ dramatically

improves unitarity up to about 2 GeV, R_0^0 violates unitarity to a lesser extent starting around 500 MeV. (As noted in Ref. 5, the $I = J = 0$ channel is the only troublesome one.) To recover unitarity, we need a negative contribution to the real part above this point, while below this point the positive contribution is preferred by experiment. Such behavior matches with the real part of a typical resonance contribution. The resonance contribution is positive in the energy region below its mass, while it is negative in the energy region above its mass. Then I include a low mass broad scalar resonance, which has historically been denoted as the σ . The σ contribution to the real part of the amplitude component $A(s, t, u)$ is given by

$$A_\sigma(s, t, u) = \frac{\gamma_\sigma^2}{2} \frac{(s - 2m_\pi^2)^2}{M_\sigma^2 - s - iM_\sigma G'} , \quad (12)$$

where the factor $(s - 2m_\pi^2)^2$ is due to the derivative-type coupling required for chiral symmetry in Eq. (8). G' is a parameter which we introduce to regularize the propagator. It can be called a width, but it turns out to be rather large so that, after the ρ and π contributions are taken into account, the partial wave amplitude R_0^0 does not clearly display the characteristic resonant behavior. In the most general situation one might imagine that G could become complex as in Eq. (11) due to higher order in $1/N_C$ corrections. It should be noted, however, that Eq. (11) expresses nothing more than the assumption of unitarity for a *narrow* resonance and hence should not really be applied to the present broad case. A reasonable fit was found in Ref. 5 for G purely real, but not equal to G' .

A best overall fit is obtained with the parameter choices; $M_\sigma = 559$ MeV, $\gamma_\sigma = 7.8 \text{ GeV}^{-1}$ and $G' = 370$ MeV. These have been slightly fine-tuned from the values in Ref. 5 in order to obtain a better fit in the 1 GeV region. The result for the real part R_0^0 due to the inclusion of the σ contribution along with the π and ρ contributions is shown in Fig. 2. It is seen that the unitarity bound is satisfied and there is a reasonable agreement with the experimental points[13, 14] up to about 800 MeV.

Next, let us consider the 1 GeV region. Reference to Fig. 2 shows that the experimental data for R_0^0 lie considerably lower than the $\pi + \rho + \sigma$ contribution between 0.9 and 1.0 GeV and then quickly reverse sign above this point. This is caused by the existence of $f_0(980)$. As we can see easily, a naive inclusion of $f_0(980)$ does not reproduce the experimental data, since the real part of the typical resonance form gives a positive contribution in the energy region below its mass, while it gives a negative contribution in the energy region above its mass. However, we need negative contribution below 1 GeV and positive contribution above 1 GeV.




Figure 2: The solid line is the *current algebra* + $\rho + \sigma$ result for R_0^0 . The experimental points, in this and succeeding figures, are extracted from the phase shifts. ($\square$) are extracted from the data of Ref. 13 while ($\triangle$) are extracted from the data of Ref. 14. The predicted R_0^0 is small around the 1 GeV region.




Figure 3: The solid line is the *current algebra* + $\rho + \sigma + f_0(980)$ result for R_0^0 obtained by assuming the values in Table 2 for the σ and $f_0(980)$ parameters.

As we discussed around Eq. (11), the effect of the background is important in this $f_0(980)$ region. In this case the background is given by the $\pi + \rho + \sigma$ contribution. Figure 2 shows that the real part of the background is very small so that the background phase δ in Eq. (11) is expected to be roughly 90° . This background effect generates the extra minus sign in front of the $f_0(980)$ contribution, as we can see from Eq. (11). Thus the $f_0(980)$ gives a negative contribution below the resonance position and gives a positive contribution above it. This is clearly exactly what is needed to bring experiment and theory into agreement up till about 1.2 GeV.

The actual amplitude used for the calculation properly contains the effects of the pions' derivative coupling to the $f_0(980)$:

$$A_{f_0(980)}(s, t, u) = e^{2i\delta} \frac{\gamma_{f_0\pi\pi}^2}{2} \frac{(s - 2m_\pi^2)^2}{M_{f_0}^2 - s - iM_{f_0}\Gamma_{f_0}}, \quad (13)$$

where δ is a background phase parameter and the real coupling constant $\gamma_{f_0\pi\pi}$ is related to the $f_0(980) \rightarrow \pi\pi$ width by

$$\Gamma(f_0 \rightarrow \pi\pi) = \frac{3}{64\pi} \frac{\gamma_{f_0\pi\pi}^2}{M_{f_0}} (M_{f_0}^2 - 2m_\pi^2)^2 \sqrt{1 - \frac{4m_\pi^2}{M_{f_0}^2}}. \quad (14)$$

The background phase parameter δ is predicted by

$$\frac{1}{2}\sin(2\delta) \equiv \tilde{R}_0^0(s = M_{f_0}^2) , \quad (15)$$

where $\tilde{R}_0^0$ is computed as the sum of the current algebra, ρ , and sigma pieces.

A best fit of our parameters to the experimental data results in the curve shown in Fig. 3. Only the three parameters $\gamma_{f_0\pi\pi}$, G' and M_σ are essentially free. The others are restricted by experiment. Since the total width of $f_0(980)$ has a large uncertainty (40 – 400 MeV in PDG list), we also fit this. In addition we have considered the precise value of M_{f_0} to be a parameter for fitting purpose. The best fitted values are shown in Table 2 together with the predicted background phase δ and the χ^2 value. The predicted background phase is seen to be close to 90° . Note that the fitted width of the $f_0(980)$ is near the low end of the experimental range in the PDG list. The low lying sigma has a mass of around 560 MeV and a width of about 370 MeV.

$M_{f_0(980)}$	$\Gamma_{f_0(980)}$	M_σ	G'	$\gamma_{f_0\pi\pi}$	δ (deg.)	χ^2
987	64.6	559	370	7.8	85.2	2.0

Table 2: The best fitted values of the parameter together with the predicted background phase δ and the χ^2 value. The unit of $M_{f_0(980)}$, $\Gamma_{f_0(980)}$, M_σ and G' is MeV and that of $\gamma_{f_0\pi\pi}$ is GeV^{-1} .

Strictly speaking our initial assumption only entitles us to compare, as we have already done, the real part of the predicted amplitude with the real part of the amplitude deduced from experiment. Since the predicted $R_0^0(s)$ up to 1.2 GeV satisfies the unitarity bound (within the fitting error) we can calculate the imaginary part, and hence the phase shift $\delta_0^0(s)$ on the assumption that full unitarity holds. This is implemented by substituting $R_0^0(s)$ into

$$\delta_0^0(s) = \frac{\arcsin(2R_0^0(s))}{2} , \quad (16)$$




Figure 4: Estimated phase shift using the predicted real part and unitarity relation.

and resolving the discrete sign ambiguities by demanding that $\delta_0^0(s)$ be continuous and monotonically increasing (to agree with experiment). The resultant phase shift $\delta_0^0(s)$ is shown in Fig. 3. As expected, the agreement is reasonable.

4 Summary

In this talk I showed the main mechanism of the analysis done in Ref. 1: (1) motivated by the large N_C approximation to QCD, we include the resonances with masses up to an energy slightly greater than the range of interest, and use the chiral symmetry to restrict the forms of the interactions; (2) the *current algebra* + ρ contribution violates the unitarity bound around 560 MeV region but it is recovered by including the low mass broad resonance sigma[5]; (3) the $\pi + \rho + \sigma$ contribution gives an important background effect to the $f_0(980)$ contribution, i.e., the sign in front of the $f_0(980)$ contribution is reversed by the background effect. The third mechanism, which leads to a sharp dip in the $I = J = 0$ partial wave contribution to the $\pi\pi$ -scattering cross section, can be identified with the very old *Ramsauer-Townsend* effect [15] which concerned the scattering of 0.7 eV electrons on rare gas atoms. The dip occurs because the background phase of $\pi/2$ causes the phase shift to go through π (rather than $\pi/2$) at the resonance position. (Of course, the cross section is proportional to $\sum_{I,J}(2J+1)\sin^2(\delta_I^J)$.) This simple mechanism seems to be all that is required to understand the main feature of $\pi\pi$ scattering in the 1 GeV region.

The detailed analysis, which includes the effects of the inelasticity ($\pi\pi \rightarrow K\bar{K}$ channel opens at 990 MeV.) and the next group of resonances, is done in Ref. 1. The results show that those effects only fine-tune the best fitted values shown in Table 2.

Acknowledgement

I would like to thank Joe Schechter and Francesco Sannino for careful reading of this manuscript, and the organizing committee of the MRST meeting for the opportunity to present this talk.

References

- [1] M. Harada, F. Sannino, and J. Schechter, Syracuse University preprint **SU-4240-624** / **hep-ph/9511335**, (1995), to be published in Phys. Rev. D.
- [2] G. 'tHooft, Nucl. Phys. **B75**, 461 (1974); E. Witten, Nucl. Phys. **B149**, 285 (1979).

- [3] S. Weinberg, *Physica* **96A**, 327 (1979); J. Gasser and H. Leutwyler, *Ann. Phys. (N.Y.)* **158**, 142 (1984); *Nucl. Phys.* **B250**, 465 (1985).
- [4] J. Gasser and U.-G. Meißner, *Phys. Lett. B* **258**, 219 (1991).
- [5] F. Sannino and J. Schechter, *Phys. Rev. D* **52**, 96 (1995).
- [6] Particle Data Group, *Phys. Rev. D* **50**, 1173 (1994).
- [7] For example, see G. Janssen, B.C. Pearce, K. Holinde and J. Speth, *Phys. Rev. D* **52**, 2690 (1995); D. Morgan and M. Pennington, *Phys. Rev. D* **48**, 1185 (1993); N.A. Törnqvist, *Z. Phys.* **C68**, 647 (1995).
- [8] Ö. Kaymakçalan and J. Schechter, *Phys. Rev. D* **31**, 1109 (1985).
- [9] M. Bando, T. Kugo and K. Yamawaki, *Phys. Rep.* **164**, 217 (1988).
- [10] J. Schechter, A. Subbaraman, and H. Weigel, *Phys. Rev. D* **48**, 339 (1993).
- [11] M. Harada and J. Schechter, Syracuse University preprint **SU-4240-613 / hep-ph/9506473**, (1995), to be published in *Phys. Rev. D*.
- [12] J. Taylor, *Scattering Theory* (Krieger, 1987).
- [13] E.A. Alekseeva et al., *Sov. Phys. JETP* **55**, 591 (1982).
- [14] G. Grayer et al., *Nucl. Phys.* **B75**, 189 (1974).
- [15] L.I. Schiff, *Quantum Mechanics* (1955).